



Pilot Study Horses

The influence of feeding a high calcium, algae supplement on gastric ulceration in adult horses

T. Moir^{1*}, J. O'Brien², S. R. Hill³ and L. A. Waldron³

¹Seahorse Supplements Ltd, Christchurch, New Zealand

²Equvet Ltd, Palmerston North, New Zealand

³LWT Animal Nutrition Ltd, PO Box 119, Feilding, New Zealand 4740

Summary

Calcium is considered important in buffering excess stomach acid in mammals, including horses. Control of stomach acid is important in preventing the development of ulcers within the stomach lining, which, in horses, are considered to be caused by acid splashing. Algae supplements contain various minerals which are in natural form, as seen in all plant and feedstuffs. The current trial was conducted to examine if a high calcium algae supplement had any impact on gastric ulceration in horses, which may be due to buffering stomach acid, reducing the pH in a gradual manner, without resorting to medication. Ten horses, of either thoroughbred, standardbred or sport horse breed, were selected on the basis of the presence of ulcers in their stomach, as ascertained by endoscopy. The average ulceration score before algae supplementation was 2.2 ± 0.75 according to the EGUC scoring system. The horses were then maintained on their normal diet (unchanged from the initial ulcer scoring) by the owner with the addition of 40 g per day of the high calcium, algae based Maxia Complete[®] (Seahorse Supplements Ltd, Christchurch, NZ) for thirty days (T30). All horses were then re endoscoped to assess any change in ulceration score. All horses showed a significant improvement in ulcer score, with seven having a score of zero (fully healed, no evidence of further ulceration) and two with a score of one (some residual inflammation or keratinosis in areas of healed ulcers). This resulted in a mean score of 0.3 ± 0.48 ($P < 0.0001$: T0 versus T30) at the end of the study. This trial demonstrated that feeding an organic form of high calcium from algae reduced ulceration in horses.

Keywords: Horses: ulceration: calcium: algae

Introduction

Gastric ulceration has been studied in various horse populations, but has mainly been found in racehorses and sport horses (dressage, eventing, showjumping). Various studies have shown the number of horses affected by ulcers ranges between 16% in standardbreds (Bezdekova *et al.* 2005) and up to 100% in working racehorses (Murray *et al.* 1996). The adult horse's stomach is divided into two regions, the lower glandular area, where acid and mucous is secreted, and the upper, squamous region, which is paler and has no secretory structures and is not protected by mucous (Reese and Andrews 2009). pH in the stomach has been reported to be

strongly acid, with pH values ranging between 2.7 and 3.2 (Murray and Grodinsky 1989; Nadeau *et al.* 2000). Horses have evolved to consume small amounts of feed continuously for around 18 hours per day (Luthersson *et al.* 2009), and, as such, secrete acid continuously as well (Murray, 1997). If the stomach is empty for any appreciable length of time, which may occur during transport, before exercise or daily in animals offered a small number of large meals, this leaves them vulnerable to acid contact in the upper, unprotected region of the stomach. The resulting ulceration mainly occurs along the *Margo plicatus* equatorial region where the two tissue types meet.

* Corresponding author. E-mail: seahorse.supplements@xtra.co.nz

Table 1. EGUC scoring system (1999)

EGUC	Gastric mucosa
0	Intact epithelium
1	Intact mucosa: hyperkeratosis/hyperaemia (healing in progress)
2	Small lesions – single/multifocal
3	Large lesions – single/multifocal
4	Extensive lesions & deep ulceration

A standardised global scoring system for equine ulceration has been developed, as shown in Table 1 based on the research by McAllister *et al* (1997) and Equine Gastric Ulceration Council, (1999), which is based on degree of severity and numbers of lesions.

Algae contains many useful minerals which may be utilised in animal nutrition (Kandale *et al.*, 2011) and is now being increasingly harvested for use from clean coastal areas of the world as well as being farmed on a large scale. The algal product under test for this trial (Maxia Complete[®], Seahorse Supplements Ltd., Christchurch, NZ) contains high levels of minerals. Maxia Complete[®] contains 32% organic calcium (356,800 ppm) and other nutritional minerals (chloride 1,206 ppm; iodine 49 ppm; iron 745 ppm; magnesium 28,390 ppm; manganese 29 ppm; phosphorus 3,454 ppm; molybdenum 1.6 ppm; potassium 1,287 ppm; silicon 475 ppm; sodium 5,302 ppm; sulphur 3,756 ppm and zinc 2.4 ppm) that the algae have incorporated into their cell walls via normal metabolism, whereby the minerals are bound to sulphated polysaccharide structures (Barnes, 2003). Calcium acts as a buffer to stomach acids, reducing the pH of the stomach contents to prevent acid damage to the sensitive upper stomach wall and around the *Margo plicatus*. However, calcium in its inorganic form, whilst having this useful activity, can act very quickly in the stomach, thereby potentially reducing the necessary activity of acid, i.e. in promoting the activity of trypsin protease enzymes, which facilitate the primary hydrolysis of proteins (Frape, 2004). It is considered that naturally occurring calcium, such as that found in algae, is more slowly available, allowing protein digestion but preventing long term ulceration by buffering the stomach acid, once it has completed its proteolytic activities. This may be useful in animals which experience periods of time between feed ingestion, where ulceration poses a major risk. Previous research has demonstrated how high calcium forages can help buffer stomach acids for six hours or more (Andrews *et al.*, 2005), and previous work using Maxia Complete[®] in a laboratory model of the horses gut, showed that it had a significant buffering effect, controlling acidity in the foregut and

promoting correct hind gut fermentation (Moore-Colyer *et al.*, 2014). The current trial was conducted to investigate if the high natural calcium levels in algae can have the same effect in the field.

Materials and methods

Advertising for volunteer animals for this pilot study in the vicinity of Christchurch in New Zealand, resulted in 57 horses (25 sport horses, 20 standardbred and 12 thoroughbred) being put forward, which were endoscoped to assess individual levels of ulceration. Of these, ten were found to have ulcer scores (1 or more on EGUC scale) that made them suitable for taking part in the trial. This gave an ulcer rate of 17.5% in the selection population of volunteered animals. This test group comprised six males and four females with an average age of 4.5 years \pm 2.51. The division in terms of breeds within the trial group of ten horses was three thoroughbreds, six standardbreds (trotting horses) and one sporthorse (mixed breed). Initial scores on the standard EGUC scale averaged 2.2 \pm 0.75. After initial scoring and identification of suitable candidates (T0), the selected horses were fed their existing diets (no change – to maintain existing stomach conditions under which ulcers had established) supplemented with 40 g/d of Maxia Complete[®] (Seahorse Supplements Ltd, Christchurch, New Zealand) algae supplement, which was top dressed and mixed into each horses daily complete feed. The ulceration scores obtained from each horse at T0 were used as the baseline, and a further scoping was undertaken after the horses had been treated for 30 days (T30) to identify changes in EGUC scoring. Ulcer scoring was conducted after 16 hours of feed and water removal according to the methods recommended by Gordon (2010) which has been previously used in published work (Stowers *et al.*, 2013). The endoscopy was carried out by Mr John O'Brien BVSc. MANZCVSc. Data was analysed using T0 as the baseline point comparing the change at the T30 time point, with means and standard deviations reported. Statistics were analysed using Unistat (Release 5.5, London UK) by ANOVA in the GLM procedure of the program.

Results

All horses were healthy and maintained good body condition score throughout the 30 day trial period. The results are shown in Table 2 below.

All horses had improved ulceration scores 30 days of feeding the high calcium Maxia Complete[®] algal supplement, with seven out of ten having no signs of ulcers and

Table 2. Trial animal demographics and endoscopy results before (T0) and after (T30) feeding Maxia Complete algal supplement to adult horses with gastric ulcers

Horse No.	Age	Sex	Breed	scoping d0	Scoping d30
1	2	Male	Thoroughbred	1	0
2	3	Male	Standardbred	1	0
3	5	Female	Thoroughbred	2	0
4	4	Female	Standardbred	2	0
5	11	Male	Sporhorse	2	0
6	5	Male	Standardbred	2.5	0
7	3	Male	Standardbred	2	1
8	3	Female	Standardbred	3	1
9	5	Male	Standardbred	3	0
10	4	Female	Thoroughbred	2	1
Mean	4.5			2.2	0.3
SD	2.51			0.75	0.48

the remaining three showing improvement to low levels of inflammation and keratinosis, most likely indicative of healing. From this initial pilot study, even though animal numbers were low and their background diets varied, the data showed that top dressing a high calcium, algal based supplement such as Maxia Complete[®], significantly ($P < 0.0001$) reduced damage to the sensitive upper non-secretory part of the equine stomach, and thus reduced ulceration scores over a 30 day period.

Conclusions

In a small group of horses with ulceration, where diets and management were varied, feeding a natural source of calcium from algae via Maxia Complete[®] reduced the acid damage to the stomach mucosa over a 30 day period and resulted in reduced ulceration scores. This agrees with other research which has shown that high calcium forages (Andrews *et al.*, 2005) and other algal products tested *in vitro* have had a similar effect (Moore-Colyer *et al.*, 2014). The role and importance of the form of calcium, i.e. as a natural form rather than inorganic, may be important in the prevention of ulceration without affecting digestive functions in the stomach. Further work is warranted to compare this activity to inorganic calcium compounds, investigate efficacy in ulceration control and subsequent impact on protein digestion.

Acknowledgements

This pilot study was sponsored by Seahorse Supplements Ltd, Christchurch, NZ

Declaration of interest

T. Moir is the director of Seahorse Supplements Ltd, Christchurch, NZ

References

- Andrews F.M., Buchanan B.R., Elliot S.B., Clariday N.A., and Edwards L.H. (2005). Gastric ulcers in horses. *Journal of Animal Science*, **83**(13_suppl), E18–E21.
- Barnes H. (2003). *Oceanography and marine biology: A review*. CRC Press, Boca Katon, Florida, USA.
- Bezděková B., Jahn P., Vyskočil M., and Plachý J. (2005). Prevalence of equine gastric ulceration in Standardbred racehorses in Czech Republic. *Acta Veterinaria Brno*, **74**(1), 59–65.
- Equine Gastric Ulceration Council (2009). Standardised scoring for ulceration in horses.
- Gordon S. (2010). Personal communication.
- Frape D. (2008). *Equine nutrition and feeding*. John Wiley & Sons.
- Kandale A., Meena A.K., Rao M.M., Panda P., and Babu R. (2011). Marine algae: An introduction, food value and medicinal uses. Report of the Food and Agriculture Organisation of the United Nations.
- Luthersson N., Hou Nielsen K., Harris P., and Parkin T.D.H. (2009). Risk factors associated with Equine Gastric Ulceration Syndrome (EGUS) in 201 horses in Denmark. *Equine Veterinary Journal*, **41**(7), 625–630.
- MacAllister C.G., Andrews F.M., Deegan E., Ruoff W., and Olovson S.G. (1997). A scoring system for gastric ulcers in the horse. *Equine Veterinary Journal*, **29**, 430–433.
- Moore-Colyer M., O’Gorman D.M., and Wakefield K. (2014). An *in vitro* investigation into the effects of a marine-derived, multiminerall supplement in simulated equine stomach and hindgut environments. *Journal of Equine Veterinary Science*, **34**(3), 391–397.
- Murray M.J. (1997). Suppression of gastric acidity in horses. *Journal of the American Veterinary Medical Association*, **211**, 37–40.
- Murray M.J. and Grodinsky C. (1989). Regional gastric pH measurement in horses and foals. *Equine Veterinary Journal*, **7**(supplement), 73–76.
- Murray M.J., Schusser G., Pipers F.S., and Gross S.J. (1996). Factors associated with gastric lesions in Thoroughbred racehorses. *Equine Veterinary Journal*, **28**(5), 368–374.
- Nadeau J.A., Andrews F.M., Mathew A.G., Argenzio R.A., Blackford J.T., Sohtell M., and Saxton A.M. (2000). Evaluation of diet as a cause of gastric ulcers in horses. *American Journal of Veterinary Research*, **61**, 784–790.
- Reese R.E. and Andrews F.M. (2009). Nutrition and dietary management of equine gastric ulcer syndrome. *Veterinary Clinics of North America: Equine Practice*, **25**(1), 79–92.
- Stowers N.L., Waldron L.A., Pryor I.D., Hill S.R. and O’Brien J. (2013). Modified bio fermentation lucerne feeds for gastric ulceration in horses. *Journal of Applied Animal Nutrition*, **1**: p1–7.